### ZH->ee bb

James Heinmiller 12MAY05 Higgs Worshop

#### Outline

Dataset
Event Selection
Areas of improvement
Electron distributions
Z distributions
Jet distributions
Future plans

#### **Datasets**

Data

Z->tau+tau

Pass 2 EM1TRK skim 336.8 1/pb includes to the end of V12

403,000

MC – Pass 2 400,000 events Z->ee +X Zjj->ee+jj291,000 Zb - ee + b101,750 Zbb->ee +bb 98,000 tt->21 + bb49,500 Wjj->e+jj133,776 ZZ->inclusive 53,500 ZH->ee + bb mass=125 5,000

Area of improvement - move to V13 dataset

All Athenalized using p16-br-03 jes 5.3

#### **Event Selection**

#### Data

Remove events with bad luminosity block numbers – Cal,CFT,Jet/Met, and SMT

Remove events with bad run numbers

Remove duplicate events

Individual EM object must pass L1&L2&L3 trigger

E1\_SH30 | E1\_SHT20 | E2\_SHT20 | E3\_SHT20

E1\_SH30 | E1\_SHT20 | E2\_SHT20

E1\_SH30 | E1\_SHT20

E1\_SHT20

EM\_HI\_SH | EM\_HI\_2EM5\_SH

EM\_HI\_SH

EM\_HI

EM\_MX\_SH

EM\_MX

Primary Vertex < 60.0 cm

For MC - only Primary Vertex<60.0cm

#### Event Selection continued

#### EM objects

#### Jet objects

| detector eta | <= 2.5 | 0.05 < emf < 0.95 | hotf < 10 | n90>1

chf < 0.40

Level 1 confirmation > 0.40 CC and FWD > 0.20 ICR Pt>20

 $\Delta R > 0.40$  between jet and electron from the Z

#### Brief selection cuts and Scale Factors

All distributions are 75.0 < Z inv mass < 105 GeV and 2 jets

#### MC corrections

#### Electrons

Pt smeared

SF for pT, eta, and phi from electron reco eff

SF for track eff

Trigger efficiency

#### <u>Jets</u>

Additional smear to data resolution

SF from jet reco eff

Have not applied SF for taggability

Have not applied SF for jlip b tagger

### Areas of Improvement - Electrons

We could add the likelihood cut.

Currently are only using electrons in the central region | detector eta | < 1.1 We could expand to allow one EM object with | detector eta | < 2.5 or both out in the EC.

Electrons	Z+ N jets	Z+ 2 jets	Z+2 taggable	Z+1 jlip 4%	Z+2jlip 4%
CC only	13907	219	156	27	2
CC+EC	26973	379	269	45	3
Increase	1.94	1.73	1.72	1.67	1.5

#### Concerns

Moving out to the EC we should modify the selection cuts – Hmatrix8

Recalculate the Trigger and EM ID\*Reco Eff

Other possibilities of gaining more statistics Loosening the Hmatrix7 cut


## Areas of Improvement – Jets

Low jet reconstruction efficiency until 40 GeV

Straight Eff with Scale Factor - Central


P17 will have 6.0 Min ET which will raise this efficiency


Improved jet resolutions and JES will raise the efficiency




#### Jet Scale Factor










## B jet resolutions

ZH->ee\_bb higgs mass = 125 GeV



b radiation – the peak has shifted to ~ 104 GeV with a large width Jet Cone of 0.70 would pick up more energy

## Cut Flow


Electrons	Z+ 2 jets	Z+2 taggable	Z+1 jlip 1%	Z+1 jlip 4%	Z+2jlip 4%
Z bb					
W bb					
П					
ZZ					
Zb					
Zjj					
ZH – 125					
Expected MC					
Data	219	156		27	2

## Next steps


Finish jet reco\*id note
Repeat Marc's analysis to estable a baseline
Expand to forward electrons
Measure Z+2 jet cross sections
Z+1 b tagged cross section
compare to Z+ b jet / Z+ jet analysis
Z+2 b tagged cross section
ZH limit

# Support Slides


#### Electrons' Eta




## Electrons' phi




## Electrons' pT



## Z Inv Mass and pT



## Z Rapidity, eta, and phi




#### Jet Distributions


Removed b mesons and baryons from Zjj+eejj

included zb+eeb and zbb+eebb


Jet pt




### Jet Eta



### Jet Phi



## Delta R between the two leading jets



### **Invariant Mass**

